

# Reduction of a bi-Hamiltonian hierarchy on $T^*\mathbf{U}(n)$ to spin Ruijsenaars–Sutherland models

László Fehér, University of Szeged and Wigner RCP, Budapest

Consider the following hierarchy of evolution equations:

$$\dot{Q}_j = (iL^k Q)_{jj}, \quad \dot{L} = [\mathcal{R}(Q)(iL^k), L], \quad \text{for } (Q, L) \in \mathbb{T}_{\text{reg}}^n \times i\mathfrak{u}(n), \quad \forall k \in \mathbb{N}.$$

$L$  is an  $n \times n$  Hermitian matrix,  $Q \equiv \text{diag}(Q_1, Q_2, \dots, Q_n)$  is a diagonal unitary matrix, and  $\mathcal{R}(Q)$  is the dynamical  $r$ -matrix given below.

There is a gauge freedom in this system:

$$(Q, L) \iff (\eta Q \eta^{-1}, \eta L \eta^{-1}) \quad \forall \eta \in \mathcal{N}(n) := N_{\mathbb{T}^n}(\mathbf{U}(n)).$$

The evolutional derivations of gauge invariant ‘observables’ commute due to the CDYBE satisfied by the dynamical  $r$ -matrix:  $\mathcal{R}(Q) := 0$  on the Cartan subalgebra  $\mathfrak{gl}(n, \mathbb{C})_0 < \mathfrak{gl}(n, \mathbb{C})$  and

$$\mathcal{R}(Q) := \frac{1}{2}(\text{Ad}_Q + \text{id})(\text{Ad}_Q - \text{id})^{-1} \text{ on } \mathfrak{gl}(n, \mathbb{C})_{\perp}, \quad (\text{Ad}_Q(X) := QXQ^{-1}).$$

Plan: First, I exhibit a bi-Hamiltonian structure for this system. Then, if time permits, I shall explain why I call it ‘spin Ruijsenaars–Sutherland hierarchy’. For details, see [arXiv:1908.02467](https://arxiv.org/abs/1908.02467) [math-ph]

## Recall celebrated exactly solvable many-body models

Trigonometric Sutherland system:

$$H_{\text{Suth}} = \frac{1}{2} \sum_{k=1}^n p_k^2 + \frac{1}{2} \sum_{j \neq k} \frac{x^2}{\sin^2(q_k - q_j)}$$

Trigonometric Ruijsenaars–Schneider system:

$$H_{\text{RS}} = \sum_{k=1}^n (\cosh p_k) \prod_{j \neq k} \left[ 1 + \frac{x^2}{\sin^2(q_k - q_j)} \right]^{\frac{1}{2}}$$

Light-cone Hamiltonians of the RS system:

$$H_{\pm} = \sum_{k=1}^n e^{\pm p_k} \prod_{j \neq k} \left[ 1 + \frac{x^2}{\sin^2(q_k - q_j)} \right]^{\frac{1}{2}}$$

Describe integrable interactions of  $n$  points moving on the circle.

Generalize rational Calogero–Moser model of points on the real line:

$$H_{\text{CM}} = \frac{1}{2} \sum_{k=1}^n p_k^2 + \frac{1}{2} \sum_{j \neq k} \frac{x^2}{(q_k - q_j)^2}$$

**Bi-Hamiltonian hierarchy on  $T^*\mathbf{U}(n)$ :** We start with the manifold

$$\mathfrak{M} := \mathbf{U}(n) \times \mathfrak{iu}(n) := \{(g, L) \mid g \in \mathbf{U}(n), L \in \mathfrak{iu}(n)\}.$$

We use the real Lie algebra  $\mathfrak{gl}(n, \mathbb{C})$ , equipped with the bilinear form

$$\langle X, Y \rangle := \Im \operatorname{tr}(XY), \quad \forall X, Y \in \mathfrak{gl}(n, \mathbb{C}),$$

and the real vector space decomposition (Manin triple)

$$\mathfrak{gl}(n, \mathbb{C}) = \mathfrak{u}(n) + \mathfrak{b}(n)$$

$$\text{with } \mathfrak{b}(n) := \operatorname{span}_{\mathbb{R}}\{E_{jj}, E_{kl}, iE_{kl} \mid 1 \leq j \leq n, 1 \leq k < l \leq n\}.$$

This gives the decomposition  $X = X_{\mathfrak{u}(n)} + X_{\mathfrak{b}(n)}$  for every  $X \in \mathfrak{gl}(n, \mathbb{C})$ .

For a real function  $F \in C^\infty(\mathfrak{M})$ , the derivatives

$$D_1 F, D'_1 F \in C^\infty(\mathfrak{M}, \mathfrak{b}(n)), \quad d_2 F \in C^\infty(\mathfrak{M}, \mathfrak{u}(n)) \quad \text{are defined by}$$

$$\left. \frac{d}{dt} \right|_{t=0} F(e^{tX} g e^{tX'}, L + tY) = \langle D_1 F(g, L), X \rangle + \langle D'_1 F(g, L), X' \rangle + \langle d_2 F(g, L), Y \rangle$$

for all  $X, X' \in \mathfrak{u}(n)$  and  $Y \in \mathfrak{iu}(n)$ .

**Proposition 1.** *The following formulas define two Poisson brackets on  $C^\infty(\mathfrak{M}, \mathbb{R})$ :*

$$\{F, H\}_1(g, L) = \langle D_1 F, d_2 H \rangle - \langle D_1 H, d_2 F \rangle + \langle L, [d_2 F, d_2 H] \rangle,$$

*and*

$$\begin{aligned} \{F, H\}_2(g, L) = & \langle D_1 F, L d_2 H \rangle - \langle D_1 H, L d_2 F \rangle \\ & + 2 \langle L d_2 F, (L d_2 H)_{\mathfrak{u}(n)} \rangle - \frac{1}{2} \langle D'_1 F, g^{-1} (D_1 H) g \rangle, \end{aligned}$$

*where the derivatives are taken at the point  $(g, L)$ .*

Remark: The first bracket is the canonical Poisson bracket of the cotangent bundle, expressed in terms of right-trivialization and taking  $\mathfrak{iu}(n)$  and  $\mathfrak{b}(n)$  as models of  $\mathfrak{u}(n)^*$ . The restriction of the second bracket to the open submanifold  $U(n) \times \exp(\mathfrak{iu}(n)) \subset \mathfrak{M}$  is a convenient multiple of Semenov-Tian-Shansky's non-degenerate Poisson bracket on the Heisenberg double of the standard Poisson-Lie group  $U(n)$ .

[Remark:  $GL(n, \mathbb{C}) \ni K = b_L g_R^{-1} = g_L b_R^{-1} \mapsto (g_R, b_R b_R^\dagger) \in U(n) \times \exp(\mathfrak{iu}(n))$ ]

Introduce the vector field  $\mathcal{D}$  on  $\mathfrak{M}$  that acts as the following derivation of the ‘coordinate functions’

$$\mathcal{D}[g_{ij}] := 0, \quad \mathcal{D}[L_{ij}] := \delta_{ij}.$$

Its flow through  $(g(0), L(0))$  reads  $(g(t), L(t)) = (g(0), L(0) + t\mathbf{1}_n)$ .

**Proposition 2.** *For  $F \in C^\infty(\mathfrak{M})$ , let  $\mathcal{D}[F]$  denote the derivative along the vector field  $\mathcal{D}$ . The Poisson brackets on  $C^\infty(\mathfrak{M})$  satisfy*

$$\{F, H\}_1 = \{F, H\}_2^{\mathcal{D}} \equiv \mathcal{D}[\{F, H\}_2] - \{\mathcal{D}[F], H\}_2 - \{F, \mathcal{D}[H]\}_2,$$

$$\{F, H\}_1^{\mathcal{D}} \equiv \mathcal{D}[\{F, H\}_1] - \{\mathcal{D}[F], H\}_1 - \{F, \mathcal{D}[H]\}_1 = 0,$$

and thus they define an exact bi-Hamiltonian structure.

The Hamiltonians  $H_k(g, L) := \frac{1}{k} \text{tr}(L^k)$  ( $\forall k \in \mathbb{N}$ ) satisfy

$$\{F, H_k\}_2 = \{F, H_{k+1}\}_1$$

and induce the bi-Hamiltonian ‘free flows’

$$(g(t), L(t)) = \left( \exp(itL(0)^k)g(0), L(0) \right).$$

Consider the following action of the group  $U(n)$  on  $\mathfrak{M}$ :

$$A_\eta(g, L) = (\eta g \eta^{-1}, \eta L \eta^{-1}), \quad \forall \eta \in U(n), (g, L) \in \mathfrak{M}.$$

One can show that the ring of invariant functions is closed under both Poisson brackets.

**Lemma 3.** *The Poisson brackets  $\{ , \}_1$  and  $\{ , \}_2$  on  $C^\infty(\mathfrak{M})$  induce two compatible Poisson brackets on  $C^\infty(\mathfrak{M})^{U(n)}$ .*

Noting that  $H_k$  is  $U(n)$  invariant, we can perform Poisson reduction, i.e., take quotient by  $U(n)$ . From now on we restrict our attention to the dense open subset

$$\mathfrak{M}_{\text{reg}} := U(n)_{\text{reg}} \times \mathfrak{iu}(n).$$

Every  $U(n)$  orbit in  $\mathfrak{M}_{\text{reg}}$  contains representatives in the submanifold

$$S := \mathbb{T}_{\text{reg}}^n \times \mathfrak{iu}(n) \subset \mathfrak{M}_{\text{reg}} \quad (\text{'gauge slice'})$$

and this submanifold is preserved by the action of the normalizer,  $\mathcal{N}(n)$ , of  $\mathbb{T}^n$  in  $U(n)$ . The embedding  $\iota : \mathbb{T}_{\text{reg}}^n \times \mathfrak{iu}(n) \rightarrow \mathfrak{M}_{\text{reg}}$  yields the identification

$$C^\infty(\mathfrak{M}_{\text{reg}})^{U(n)} \simeq C^\infty(\mathbb{T}_{\text{reg}}^n \times \mathfrak{iu}(n))^{\mathcal{N}(n)} \quad (\text{'restricted invariants'})$$

We obtain the reduced Poisson algebras  $(C^\infty(\mathbb{T}_{\text{reg}}^n \times \mathfrak{iu}(n))^{\mathcal{N}(n)}, \{ , \}_i^{\text{red}})$ :

$$\{F \circ \iota, H \circ \iota\}_i^{\text{red}} := \{F, H\}_i \circ \iota \quad \text{for } F, H \in C^\infty(\mathfrak{M}_{\text{reg}})^{\cup(n)}, i = 1, 2.$$

Using  $\mathcal{R}(Q) \in \text{End}(\mathfrak{gl}(n, \mathbb{C}))$ , introduce

$$[X, Y]_{\mathcal{R}(Q)} := [\mathcal{R}(Q)X, Y] + [X, \mathcal{R}(Q)Y], \quad \forall X, Y \in \mathfrak{gl}(n, \mathbb{C}).$$

For any  $f \in C^\infty(\mathbb{T}_{\text{reg}}^n \times \mathfrak{iu}(n))$ , we have the  $\mathfrak{b}(n)_0$ -valued derivative  $D_1 f$  and the  $\mathfrak{u}(n)$ -valued derivative  $d_2 f$ :

$$\langle D_1 f(Q, L), X \rangle + \langle d_2 f(Q, L), Y \rangle = \left. \frac{d}{dt} \right|_{t=0} f(e^{tX} Q, L + tY).$$

**Theorem 4.** For  $f, h \in C^\infty(\mathbb{T}_{\text{reg}}^n \times \mathfrak{iu}(n))^{\mathcal{N}(n)}$ , the reduced Poisson brackets obey the explicit formulas

$$\{f, h\}_1^{\text{red}}(Q, L) = \langle D_1 f, d_2 h \rangle - \langle D_1 h, d_2 f \rangle + \langle L, [d_2 f, d_2 h]_{\mathcal{R}(Q)} \rangle,$$

and

$$\{f, h\}_2^{\text{red}}(Q, L) = \langle D_1 f, L d_2 h \rangle - \langle D_1 h, L d_2 f \rangle + 2 \langle L d_2 f, \mathcal{R}(Q)(L d_2 h) \rangle,$$

where the derivatives are evaluated at the point  $(Q, L)$ .

**Theorem 5.** *The bi-Hamiltonian vector field  $V_k$  on  $\mathfrak{M}$ , given by*

$$V_k[F] =: \{F, H_k\}_2 = \{F, H_{k+1}\}_1, \quad k \in \mathbb{N},$$

*induces a derivation of  $C^\infty(\mathbb{T}_{\text{reg}}^n \times \mathfrak{iu}(n))^{\mathcal{N}(n)}$ . Up to infinitesimal gauge transformations, this is given by the vector field  $W_k$  on  $\mathbb{T}_{\text{reg}}^n \times \mathfrak{iu}(n)$  that satisfies*

$$\dot{Q}Q^{-1} := W_k[Q]Q^{-1} = (iL^k)_{\text{diag}}, \quad \dot{L} := W_k[L] = [\mathcal{R}(Q)(iL^k), L].$$

*As derivations of  $\mathcal{N}(n)$ -invariant functions,  $f = F \circ \iota$  and  $h_k = H_k \circ \iota$ , these reduced evolutional derivations obey*

$$W_k[f] = \{f, h_k\}_2^{\text{red}} = \{f, h_{k+1}\}_1^{\text{red}}.$$

Summary: We have shown that Poisson reduction of the bi-Hamiltonian hierarchy of ‘free motion’ on  $\mathfrak{M} = T^*\text{U}(n)$  results in a bi-Hamiltonian hierarchy describing the time development of the gauge invariant observables forming  $C^\infty(\mathbb{T}_{\text{reg}}^n \times \mathfrak{iu}(n))^{\mathcal{N}(n)}$ . The reduced hierarchy is called ‘trigonometric spin Ruijsenaars–Sutherland hierarchy’.



**Interpretation as a spin Sutherland model (well-known):** Introduce new variables by the diffeomorphism:

$$\mathbb{T}_{\text{reg}}^n \times \mathfrak{iu}(n) \ni (Q, L) \iff (Q, p, \phi) \in \mathbb{T}_{\text{reg}}^n \times \mathfrak{iu}(n)_{\text{diag}} \times \mathfrak{iu}(n)_{\perp}$$

$$\text{using } L(Q, p, \phi) := p + (\mathcal{R}(Q) + \frac{1}{2}\text{id})(\phi).$$

The entries  $p_j$  of  $p$  and  $q_j$  in  $Q_j = e^{iq_j}$  form canonically conjugate pairs, and are combined with the Poisson algebra of the quotient

$\mathfrak{u}(n)^* //_0 \mathbb{T}^n = (\mathfrak{iu}(n)_{\perp}) / \mathbb{T}^n$ . The space of physical observables becomes

$$C^{\infty}(\mathbb{T}_{\text{reg}}^n \times \mathfrak{iu}(n)_{\text{diag}} \times \mathfrak{iu}(n)_{\perp})^{\mathcal{N}(n)},$$

and the reduced first Poisson bracket takes the form

$$\{\mathcal{F}, \mathcal{H}\}_1^{\text{red}}(Q, p, \phi) = \langle D_Q \mathcal{F}, d_p \mathcal{H} \rangle - \langle D_Q \mathcal{H}, d_p \mathcal{F} \rangle + \langle \phi, [d_{\phi} \mathcal{F}, d_{\phi} \mathcal{H}] \rangle.$$

In these variables, we get the standard spin Sutherland Hamiltonian

$$\mathcal{H}_2(Q, p, \phi) := \frac{1}{2}(L(Q, p, \phi))^2 = \frac{1}{2} \sum_i p_i^2 + \frac{1}{8} \sum_{i \neq j} \frac{|\phi_{ij}|^2}{\sin^2 \frac{q_i - q_j}{2}}.$$

The spin variable  $\phi$  can be restricted by fixing the values of the Casimir functions  $C_i \in C^{\infty}(\mathfrak{u}(n)^*)^{\mathfrak{U}(n)}$ , and a special choice gives the spinless Sutherland model.

**Interpretation as a spin Ruijsenaars model:** Restrict attention to

$$\mathbb{T}_{\text{reg}}^n \times \exp(\mathfrak{iu}(n)) \subset \mathbb{T}_{\text{reg}}^n \times \mathfrak{iu}(n),$$

where  $L$  can be uniquely written in the form

$$L = e^p b_+ (b_+)^{\dagger} e^p \quad \text{with} \quad p \in \mathfrak{b}(n)_0, \quad b_+ \in \exp(\mathfrak{b}(n)_+) =: \mathbf{B}(n)_+.$$

Then consider the invertible change of variables

$$(Q, L) \longleftrightarrow (Q, p, b_+) \longleftrightarrow (Q, p, \lambda(Q, b_+)) \quad \text{with} \quad \lambda(Q, b_+) = b_+^{-1} Q^{-1} b_+ Q.$$

$\lambda$  varies freely in the triangular nilpotent subgroup  $\mathbf{B}(n)_+ < \mathbf{B}(n)$ . This gives the identification

$$C^\infty \left( \mathbb{T}_{\text{reg}}^n \times \exp(\mathfrak{iu}(n)) \right)^{\mathcal{N}(n)} \longleftrightarrow C^\infty \left( \mathbb{T}_{\text{reg}}^n \times \mathfrak{b}(n)_0 \times \mathbf{B}(n)_+ \right)^{\mathcal{N}(n)}.$$

*For  $\mathcal{F}, \mathcal{H} \in C^\infty(\mathbb{T}_{\text{reg}}^n \times \mathfrak{b}(n)_0 \times \mathbf{B}(n)_+)^{\mathcal{N}(n)}$ , the change of variables leads to the ‘decoupled form’ of the second Poisson bracket:*

$$2\{\mathcal{F}, \mathcal{H}\}_2^{\text{red}}(Q, p, \lambda) = \langle D_Q \mathcal{F}, d_p \mathcal{H} \rangle - \langle D_Q \mathcal{H}, d_p \mathcal{F} \rangle + \langle D'_\lambda \mathcal{F}, \lambda^{-1} (D_\lambda \mathcal{H}) \lambda \rangle.$$

The last term encodes the natural Poisson bracket on  $\mathbf{B}(n) //_0 \mathbb{T}^n$ , which is the Poisson-Lie analogue of  $\mathfrak{u}(n)^* //_0 \mathbb{T}^n$ .

In terms of these variables, the main Hamiltonian  $\text{tr}(L)$  has the form

$$\text{tr}(L) = \sum_{i=1}^n e^{2p_i} V_i(Q, \lambda) \quad \text{with} \quad V_i(Q, \lambda) = \left( b_+(Q, \lambda) b_+(Q, \lambda)^\dagger \right)_{ii},$$

and thus the reduced system can be interpreted as a spin RS model. The corresponding open subset of the reduced phase space is

$$\left( \mathbb{T}_{\text{reg}}^n \times \mathfrak{b}(n)_0 \times (\mathbb{B}(n)_+ / \mathbb{T}^n) \right) / S_n.$$

We obtain Poisson subspaces by restricting  $\mathbb{B}(n)_+ / \mathbb{T}^n$  to  $\mathbb{T}^n$ -reduced dressing orbits of  $U(n)$ . The dressing orbits  $\tilde{\mathcal{O}} \subset \mathbb{B}(n)$  are obtained by fixing the Casimirs,  $\mathcal{C}_i \in C^\infty(\mathbb{B}(n))^{U(n)}$ . The smallest non-trivial dressing orbit gives the standard *spinless, trigonometric* (real) RS model.

- Does the bi-Hamiltonian story generalize in a reasonable manner if we replace  $U(n)$  by an arbitrary compact simple Lie group?
- What about generalization to spin Sutherland and RS models of Gibbons–Hermsen and Krichever–Zabrodin type, and what about the elliptic case?
- Outstanding open question: **How to obtain the standard spinless, hyperbolic (real, repulsive) RS model by Hamiltonian reduction?**

## A very incomplete list of references

- D. Kazhdan, B. Kostant and S. Sternberg: *Hamiltonian group actions and dynamical systems of Calogero type*, Comm. Pure Appl. Math. **XXXI** (1978) 481-507
- J. Gibbons and T. Hermsen: *A generalisation of the Calogero–Moser system*, Physica D **11** (1984) 337-348
- M.A. Semenov-Tian-Shansky, *Dressing transformations and Poisson group actions*, Publ. RIMS **21** (1985) 1237-1260
- P.I. Etingof, I.B. Frenkel and A.A. Kirillov Jr.: *Spherical functions on affine Lie groups*, Duke Math. J. **80** (1995) 59-90
- I. Krichever and A. Zabrodin: *Spin generalization of the Ruijsenaars–Schneider model, non-abelian 2D Toda chain and representations of Sklyanin algebra*, Russian Math. Surveys **50** (1995) 1101-1150
- L.-C. Li and P. Xu: *A class of integrable spin Calogero–Moser systems*, Commun. Math. Phys. **231** (2002) 257-286
- L. F. and B.G. Puztai: *Spin Calogero models obtained from dynamical  $r$ -matrices and geodesic motion*, Nucl. Phys. B **734** (2006) 304-325
- L.F. and C. Klimčík: *Poisson-Lie generalization of the Kazhdan–Kostant–Sternberg reduction*, Lett. Math. Phys. **87** (2009) 125-138
- N. Reshetikhin: *Degenerately integrable systems*, J. Math. Sci. **213** (2016) 769-785
- L.F.: *Poisson-Lie analogues of spin Sutherland models*, arXiv:1809.01529 [math-ph]
- L.F.: *Reduction of a bi-Hamiltonian hierarchy on  $T^*U(n)$  to spin Ruijsenaars–Sutherland models*, arXiv:1908.02467 [math-ph]